

Appendix 1. Species occurrence and abundance on and off burrowing bettong warrens (n = 10) over two years. “Glenora” Station, Bourke, N.S.W.

Key: ■ = Major component of biomass and present at several sites; □ = Minor component of biomass and only present at few sites; — = Not present.

Species	On Warren		Off Warren	
	1997	1998	1997	1998
<hr/>				
Forbs				
<i>Abutilon</i> sp.	—	—	—	□
<i>Boerhavia diffusa</i>	□	■	□	□
<i>Calotis hispidula</i>	—	■	—	■
<i>Carthamus lanatus</i>	—	□	—	—
<i>Chenopodium</i> sp.	—	□	—	—
<i>Craspedia</i> sp.	—	□	—	□
<i>Rhodanthe floribunda</i>	—	■	—	■
<i>Rhodanthe</i> sp.	□	□	—	□
<i>Maireana</i> sp.	—	—	□	—
<i>Medicago</i> sp.	—	□	—	—
<i>Pimelia</i> sp.	—	—	□	—
<i>Ptilotus obovatus</i>	—	■	□	■
<i>Salsola kali</i>	□	—	—	—

<i>Sclerolaena diacantha</i>	■	■	■	■
<i>Sclerolaena quinquecuspis</i>	■	—	□	—
<i>Sida</i> sp.	□	■	□	□
<i>Solanum</i> sp.	—	□	—	□
<i>Vittadinea</i> sp.	—	□	—	—
<i>Wahlenbergia</i> sp.	—	□	—	—

---

Palatable grasses

<i>Digitaria</i> sp.	□	—	□	—
<i>Enneapogon avenaceus</i>	■	■	□	□
<i>Eragrostis</i> sp.	—	—	—	□
<i>Monachather paradoxa</i>	—	□	□	■
<i>Austrostipa</i> sp.	—	■	—	■
<i>Thyridolepis mitchelliana</i>	—	—	—	■

---

Unpalatable grasses

<i>Aristida contorta</i>	□	□	□	□
<i>Eragrostis eriopoda</i>	□	□	■	■

---